

THE UNITED STATES OF AVIORIDA

TO MILTOWHOM THESE PRESENTS SHAM COME:

Hnibersity of Georgin Research Joundation, Inc. (HGARI)

and Ilorida Agricultural Experiment Station (IAES)

MICCOLS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN EQUING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY ACCION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT, COMMON

'518W

In Testimone Morrest, I have hereunto set my hand and caused the seal of the Hunt Unriety Protestion Office to be affixed at the City of Washington, D.C. this twelfth day of September, in the year two thousand one.

Makenenon

Scretary of Agriculture

Alisti Pal M. Jaukon

Commissioner Plant Variety Protection C Agricultural Marketing S

SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTE	CTION OFFICE	1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 199		
APPLICATION FOR PLANT VARIETY PROTECTION	Application is required in order to determine if a plant variety protectificate is to be issued (7 U.S.C. 2421). Information is held confidunt if certificate is issued (7 U.S.C. 2426).			
University of Georgia Research Found (UGARF) and Florida Agricultural Exp Station (FAES)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER GA87467E24	3. VARIETY NAME 518W		
4. ADDRESS (Street and Mo. or R.F.D. Mo. Chy, State, and ZP Code, and Course Boyd Graduate Studies Research Center Room 630, D.W. Brooks Drive Athens, GA 30602-7411	er	6. TELEPHONE Shokude area code) 706-542-6512 6. FAX Shokude area code) 706-542-3837	FOR OFFICIAL USE ONLY 98.00327	
7. GENUS AND SPECIES NAME		• .	6/23/98	
Triticum aestivum	Gramineae		FLING AND EXAMINATION FEE:	
8. CROP KIND NAME (Common name)			E DATE	
Wheat, common			6/23/98	
O. IF THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATIO	N (corporation, pertnership	strocistion etc.) Comme somel	C CERTIFICATION FEE: AZ	
Corporation (UGARF) and University (FAES)		320	
Georgia (UGARF) NAME AND ADDRESS OF APPLICANT REPRESENTATIVEIS), IF ANY, TO SERVE O	ļ	November 17, 1978	E DATE S/27/01	
CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED Follow Institute. Exhibit A. Origin and Breading History of the Variety. Exhibit B. Statement of Distinctness. Exhibit C. Objective Description of the Variety. A. S. Exhibit D. Additional Description of the Variety (Optional). Exhibit E. Statement of the Basis of the Applicant's Ownership.				
Solution Sample (2,600 visible untracted seeds or, for tuber propagated variety.) X Firm and Exemination Fee (52,450) mode analysis The propagated variety.	eties verification that tissue	culture will be deposited and maintained	in an approved subfir managing of	
tva. and the Unit	led States" Bled to PVPOI			
DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY VARIE (YES 87 Yes, answer from 18 and 19 below)	CALINO III NO, DE NO NE	ASS OF CERTIFIED SEED? (See Section 6	13(a) of the Plant Variety Protection Acti	
DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO	NUMBER OF 18. IF	YES" TO ITEM 18, WHICH CLASSES OF	PRODUCTION REYOND RREEDER SEED?	
☐ YES ☐ NO	1	FOUNDATION REGISTERED		
AS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY BEEN RELEASED YES 87 'yes, "give names of countries and detail."	. USED, OFFERED FOR SA	LE OR MARKETED IN THE U.S. OR OTH	CERTIFIED	
YES 81 "yes," give names of countries and dates!	****		in coon inggr	
		•		
he applicantial declare that a viable sample of basic seed of the variety will be furnish applicable, or far a tuber propagated variety a tissue culture will be deposited in a publ	ed with application and wi	I be replenished upon request in accordan	- with the state of the state o	
A ≥ 7 × ×	a separate from the separate of	- I A DE DESCRIPTION OF THE CONTRACTOR		
te undersigned applicantis) is(are) the owner(a) of this sexually reproduced or tuber pr ction 42, and is entitled to protection under the provisions of Section 42 of the Plant	ropagated plant variety, and Variety Pretection Act.	d believels) that the variety is new, distinc	It, uniform, and stable as required in	
opticant(s) isfare) informed that false representation herein can jeopardize protection as				
TURE OF APPLICANT POWNERFAIL		APPLICANT (Owner(s))		
Please printer ofpe)	NAME (Please pr	nt or typel		
TY OR TITLE		•		
Executive Vice President 6-22-	CAPACITY OR TO	TLE	DATE	
70 mags in				

Exhibit A Origin and Breeding History of 518W

'518W ' a soft red winter wheat (Triticum aestivum L.) was cooperatively developed and released by the Georgia and Florida Agricultural Experiment Stations in 1997. 518W was derived from a three way cross in 1987: GA 80573 // 'GA-Gore' / 'Balkan'. 80573 is an experimental line from the cross, SS Tuscan V/'Blueboy'. Balkan is an European line which possesses the Solid Straw Tuscan Variant, PI 108980, was used because it possesses a solid stem and large seeds. The cultivar was developed using a modified pedigree method of breeding. was grown in the field during the 1987 season. Individual spike selections were made in the F2 thru F5 generations for leaf rust, powdery mildew, septoria nodorum and agronomic traits at Plains, 518W is the F5-derived bulk of seven F8 head rows bulked selected from 100 head rows. Breeder seed, produced in 1997, is in the F10 generation.

518W was evaluated for agronomic performance as GA 87467E24 in breeding nursery plots in 1993 (1 rep at two locations) and 1994 (4 rep at four locations), in state trials at five locations in 1995 thru 1997, and in the Uniform Southern Soft Red Winter Wheat Nursery at about 29 locations in 1997.

518W is an medium maturing, awnless, white chaffed, short statured (88 cm) at maturity, apically awnletted and has moderate straw strength. During the 3 yr (five locations yr-1 in Georgia, 518W and GA-Gore yielded an average of 3808 and 3806 kg ha-1, respectively. It matures on average 1 day earlier than GA-Gore. Milling and baking quality characteristics of 518W are rated as adequate for soft red winter wheat by the USDA-Soft Wheat Quality Laboratory, Wooster, OH.

518W is susceptible to all biotypes of Hessian fly (Mayetiola destructor (Say)) in Georgia. It is resistant to current races of leaf rust caused by <u>Puccinia recondita</u> (Roberge ex Desmaz), and moderately resistance to powdery mildew (<u>Erysiphe graminis</u> DC. f. sp. <u>tritici</u> Em. Marchal) in the Southeast.

518W has remained uniform and stable in composition through five generation of selfing in the field. Variants are limited to less than 1% and may include slightly tall late and awned plants.

Revised Exhibit B

Novelty Statement 518W

518W is a soft red winter wheat which is uniquely different but is most similar in appearance to one its parent GA-Gore. However, 518W has a clavate head shape and crooked peduncle whereas GA-Gore has a oblong head shape and straight peduncle.

518W is susceptible in the field to biotypes O and M for Hessian fly while GA-Gore is resistant (attached table). 518W is susceptible to all biotypes tested in the greenhouse, USDA-ARS Hessian Fly Laboratory, Purdue University.

Hessian Fly infestations of GA 87467 and check cultivars 1996 and 1997 at Plains, Georgia.

Entry	% Infestedstems		Immatures per_stem
	1996	1997	1996 1997
GA 87467	24a	16a	0.37a 0.28a
Gore	1b	1b	0.00b 0.01b
C9835	0b	0b	0.00b 0.00b
PIO 2684	4b	1b	0.04b 0.01b
Lsd (0.05)	12	9	0.069 0.16

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE COMMODITIES SCIENTIFIC SUPPORT DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (TRITICUM SPP.)

INSTRUCTION.	
University of Georgia Research Foundation, Inc.	FOR OFFICIAL USE ONLY
ADDRESS (Street and No. or R. F.D. No., City, State, and ZIP Code)	PVPO NUMBER Q OAC 277
NOUNCES (Client will fine or fur to, story, orace, and ZIP Code)	980327
	DESIGNATION
	518W
Place the appropriate number that describes the varietal character of this variety in Place a zero in first box (c.s. 0 8 9 or 0 9) when number is either 99 or less	the boxes below. or 9 or less.
1. KIND:	
	DULARD 7 = CLUB
2. TYPE: 2 I = SPRING 2 = WINTER 3 = OTHER (Specify) 1 = SOFT 2 = HARD	3 = OTHER (Specify)
2 1= WHITE 2=RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
· · · · · · · · · · · · · · · · · · ·	ST FLOWERING
4. MATURITY (SOX Flowering):	
0 1 NO. OF DAYS EARLIER THAN 7 1 = ARTHU	R 2 = scout 3 = chris
NO. OF DAYS LATER THAN	5 = NUGAINES 6 = LEEDS
5. PLAKT HEIGHT (From sell level to top of head):	
0 8 8 cm. High	
CM. TALLER THAN	-
0 5 CM. SHORTER THAN	R 2 = SCOUT 3 = CHRIS 7 = GA. 5 = NUGAINES 6 = LEEDS Gore
PLANT COLOR AT BOOTING (See reverse): 7. ANTHER COLOR	<u> </u>
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN 1 1 = YELLOW	2 * PURPLE
STEM:	
1 Anthocyanin: I = ABSENT 2 = PRESENT 2 Vaxy bloom:	I = ABSENT 2 = PRESENT
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT 1 Internodes: 1	≖ HOLLOW 2 × SOLID
	ERNODE LENGTH BETWEEN FLAG LEAF AF BELOW
AURICLES:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT 1 Hairiness: 1 =	ABSENT 2 = PRESENT
. LEAF:	
Flag leaf at = ERECT 2 = RECURVED 1 Flag leaf: =	HOT TWISTED 2 TWISTED
1 Haira of fires leaf sheethy 1 - anceys 2 - one cour	flag leaf obeath: = ABSENT 2 = PRESENT
1 1 MM. LEAF WIDTH (First leaf below flag loop) 2 0 CM. LEAF	F LENGTH (First leaf below flag leaf):
RM LMGS 470-6 (6-82) (Formerly Form LPGS 470-6 (3-79), which may be used)	

Density: 1 = LAX	2 = DENSE	3 Shape: 1 = TAPI 4 = OTHI	ERING 7 = STRAP 3 = CLAVATE ER (Specify)
	INLESS 2 = APICALLY AWALETED		(EO
1 Color at maturity: 5	= WHITE 2 = YELLOW 3 = PINK = BROWN 6 = BLACK 7 = OTE	4 = RED IER (Specily):	
0 6 CM. LENGTH.		1 1 MM. WIOTH	
1 2 1	TY: (CA. 7 mm.) 2 = MEDIUM (CA. 8 mm.) (CA. 9 mm.)	3 = WIDE	DW (CA. 3 mm.) 2 = MEDIUM (CA. 3.5 mm.)
Shoulder I = WANT shape: 4 = SQUA	ing $2 = OBLIQUE$ $3 = ROUNDED$ RE $5 = ELEVATED$ $6 = APICULATE$	2 Bcak: 1 = OBTUS	E 2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLOR	:	14. SEEDLING ANTHOO	YANIN:
I = WHITE 2 = R	ED 3 = PURPLE	1 = ABSENT	2 = PRESENT
15. JUVENILE PLANT GR	OWTH HABIT:		
2 1 = PROSTRATE	2 = SEMI-ERECT 3 = ERE	EGT	And the second s
16. SEED:			
1 Shape: 1 = OVATE	2 = OVAL 3 = ELLIPTICAL	1 Check: 1 = ROUN	DED 2 = ANGULAR
1 Brush, 1 = SHORT	2 = MEDIUM 3 = LONG	I Brush: 1 = HOT r	COLLARED 2 = COLLARED
Phenol reaction	1 = IVORY 2 = FAWN 3 = LT. BROY	•	- COLUMN - 1 - COLUMN
(See instructions):	4 = BROWN 5 = BLACK		
Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	S = OTHER (Specity)	
0 6 MM. LENGTH	0 4 MM. WIDTH	3 8 GM. PER 1000	SEEDS
17. SEED CREASE:			
± [ESS OF KERNEL WINOKA] 4	R LESS OF KERNEL 'SCOUT'
<u>-</u>	ESS OF KERNEL 'CHRIS'		R LESS OF KERNEL 'CHRIS' R LESS OF KERNEL "LEMHI"
	IS WIDE AS KERNEL "LEMHI" led, 1 = Susceptible, 2 = Resistant)	3 30 t C.	grand the state of
2 STEM RUST QFCQ	2 LEAF RUST PLMQ, MCJI	STRIPE RUST	0 LOOSE SMUT
2 POWDERY MILDEW	BUNT	OTHER (Specify)	
19. INSECT: (0 = Not Teste	d, 1 = Susceptible, 2 = Resistant)		
SAWFLY	O APHID (Bydv.)	O GREEN BUG	O CEREAL LEAF BEETLE
OTHER (Spectfy)	HESSIAN FLY	I GP I A	1 s 1 c
• .	RACES:	1 o 1 e	0 F 1 6
O. INDICATE WHICH VARIS	TY MOST CLOSELY RESEMBLES THAT S	(IIRUITTED:	
CHARACTER	NAME OF VARIETY	CHARACTER	HAME OF VARIETY
Plant tillering	GA-GORE	Seed size	GA-GORE
Leof size	GA-GORE	Seed shape	GA-CORE
Leaf color	GA-GORE /O: W ZE 1	Colcoptile elongation	
Leaf carriage	GA-GORE Z	Seedling pigmentation	2 26.95 3

INSTRUCTIONS

GENERALI. The following publications may be used as a reference aid for the and addition of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1963. A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

Exhibit D

Additional Description of 518W

518W is a common soft red winter wheat, <u>Triticum aestivum L.</u> bred and developed by the University of Georgia, Georgia Agricultural Experiment Stations and developed jointly by Jerry W. Johnson and Ron D. Barnett with the University of Florida, Florida Agricultural Experimental Station.

518W is medium maturing, high yielding, apically-awnleted and short in stature at maturity. It has excellent resistant to leaf rust, <u>Puccinia recondita</u> (Roberge ex Desmaz). 518W is moderately resistant to races of powdery mildew, (<u>Erysiphe graminis DC</u>. f. sp. <u>tritici Em</u>. Marchal) in the southeast region and has a moderate level of resistance to leaf and glume blotch, (<u>Stagonospora nodorum</u> and barley yellow dwarf virus. Information on the milling and baking quality characteristics is also included in a quality report. Additional information is presented in Tables 1-13 attached to this Exhibit.

Table 1. Average performance of GA 847467 and check cultivars in Elite Nursery, 1994.

Entry	Yield Bu/A	Test Wt. lbs/bu ₁	Lodging %2	Date Headed ₁	Height in
GA 87467	97.4a	58.4	30	4/04	33
Gore	89.4b	57.8	43	4/04	34
C 9835	93.9ab	57.6	33	4/07	31
PIO 2684	92.2a	62.6	22	4/04	34

¹ Plains and Griffin

Table 2. Average yield performance of GA 87467 and check cultivars over 2 years (1995-1996) at five locations.

			Location			
Entry	Tifton	Plains	Midville	Griffin	Calhoun	Average
GA 87467	50.2c	59.9a	68.9a	64.1a 3 ∈0 ∩	55.0a	59.6a
Gore	55.7b	58.9a	68.4a	69.0a	61.3a	62.7a
C 9835	48.3c	54.1a	60.9 a	71.5a	64.0a	59.6a
PIO 2684	66.4a	62.2a	72.1a	66.9a	63.8a	66.3a

² Plains, Griffin, and Calhoun

Table 3. Average performance of GA 87467 and check cultivars over 2 years (1995-1996).

Entry	Test Wt. 1bs/bu	Lodging %	Date Headed	Height in
GA 87467	55.5c	19	4/14	36
Gore	57.5b	21	4/13	37
C 9835	56.0c	5	4/15	34
PIO 2684	59.5a	12	4/14	37

Table 4. Average performance of GA 87467 and check cultivars over 2 years (1995-1996).

Entry	Leaf Rust¹ %	Powdery Mildew ₂ %	Hessian Fly %	BYD ₃ %	Septoria nodorum₄ %
GA 87467	0c	2	25	15	21
Gore	41b	, 1	1	10	15
Dozier	1c	.88 51111 53		5	8
C 9835	86a	10 0 0	1	10	8
PIO 2684	56ab	Ó.	2	5	10

¹ Tifton, Plains, 1995

² Tifton, Plains, Griffin, 1995

³ Tifton, 1995

⁴ Tifton, 1995 and Plains, 199

Table 5. Average yield performance of GA 87467 and check cultivars over 3 years (1995-1997) at five locations.

			Location	<u> </u>			Average	· · · · · · · · · · · · · · · · · · ·
Entry	Tifton	Plains	Midville	Griffin	Calhoun	S. GA.	N. GA.	STATE
GA 87467	48.4b	54.4a	67.1ab	67.2a	51.4c	56.6b	59.3b	57.7b
Gore	52.9b	47.5a	64.8b	66.1a	56.9b	55.0b	61.5b	57.6b
C 9835	48.5b	55.0a	60.9c	74.4a	59.1ab	54.8b	66.9a	59.6b
PI0 2684	62.2a	57.4a	69.0a	72.4a	62.4a	62.8a	67.0a	64.5a

Table 6. Average performance of GA 87467 and check cultivars over 3 years (1995-1997).

Entry	Test Wt. lbs/bu	Date Headed	Lodging %	Height in.	· · · · · · · · · · · · · · · · · · ·
GA 87467	55.5c	35° 44:753 44° 20	29	35	
Gore	56.8b	4-8	29	37	
C 9835	56.0bc (GL)	20V4-9	7	34	
PI0 2684	59.1a	4-8	11	36	

Table 7. Average performance of GA 87467 and check cultivars over 3 years (1995-1997).

	Leaf	Powdery	BYD	Septoria	
Entry	Rust %	Mildew %	%	Nodorum %	
GA 87467	0ь	1	28	22	
Gore	33a	1	24	15	
C 9835	53a	11	17	25	
P 2684	49a	0	11	10	

Table 8. Average performance of GA 87467 and check cultivars in Elite Nursery

at Plains and Griffin, 1997.								
Entry	Yield Bu/A	Test Wt. Ibs/bu	Lodging %	Date Headed	Height in			
GA 87467	78.3a	58.5	20	3/25	29			
Gore	66.0b	58.5	10	3/30	30			
C 9835	73.6a	59.1 FTM353	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	3/31	26			
PIO 2684	68.0b	59.9		3/28	30			

Table 9. Average performance of GA 87467 and check cultivars in other states performance trials and/or Uniform Southern Wheat Nursery (23 Trials)++, 1997.

Entry	Yield Bu/A	Test Wt Lbs/Bu	Date Headed	Lodging	Height	Leaf Rust	Powdery Mildew
GA 87467	75.8	56.9	4-14	16	35	5	4
C 9835	74.4	56.6	4-20	7	33	11	12
P 2643	72.3	57.1	4-17	8	31	23	4

++ States and (number of Locations) tested: Arkansas (6), Louisiana (1), Mississippi (2), North Carolina (4), South Carolina (4), Texas (1), Virginia (5).

Arkansas (Bay, Keiser, Kibler, Marianna, Rohwer, Hope); Louisiana (Baton Rouge); Mississippi (Raymond, Newton); North Carolina (Kinston, Bertie, Lendir, Washington); South Carolina (Simpson-early and late planting, Florence, Blacksville); Texas (Overton); Virginia (Blacksburg, Holland, Painter, Warsaw, Orange).

Table 10. Average performance of GA 87467 and check cultivars in other states performance trials and/or Uniform Southern Wheat Nursery (23 Trials)++, 1997.

Location								
Entry	AR	LA	MS	SC	NC	VA		
GA 87467	81.3	57.5	30.1	95.3	59.0	93.2		
C 9835	79.5	49.4	48.0	99.3	62.5	78.2		
PI0 2643	82.7	42.8	53.5	87.0	60.8	77.2		

Tablell.

Hessian infestations in entries of the State Wheat Variety Trial at Griffin, GA in spring of 1997.

F4.	% Infested	Immatures
Entry	stems	per stem
Clemens	80.0 *	3.07 *
Coker 9803	60.0 *	1.67 *
Jackson	58.7 *	1.71 *
GA-Dozier	57.3 *	1.25 *
Coker L910097	57.3 *	1.44 *
VA 93-52-60	54.7 *	1.16 *
Featherstone 520	53.3 *	1.67 *
Hunter	48.0 *	1.28 *
Coker 9134	42.7 *	* 19.0
Clark	42.7 *	0.99 *
Agripro 92D+4413	42.7 *	0.95 *
Kucharek-A28	37.3 *	1.03 *
GA 87467-E24	34.7 *	0.91 *
GA 891283-LE18	33.3 *	0.97 *
Mason	33.3 *	1.07 *
Agratech FFR 522W	32.0 *	0.65 *
Clemson 201	30.7 *	1.08 *
GA Gore	29.3 *	0.52
ladison	29.3 *	0.52
lorida 304	29.3 *	0.51
ioneer 2643	28.0 *	0.53
турее	26.7 *	0.59 *
ioneer 2628	16.0	0.24
ioneer 2580	16.0	0.33
ickory	12.0	0.20
oker 9835	10.7	0.25
lorey	10.7	0.21
A Fleming (90078-I)	9.3	0.27
A 87105-E43	8.0	0.20
oneer 2684	8.0	0.13
A 881186-E48	5.3	
A 871339-E18	5.3 5.3	0.08
oker 9663	5.3	0.08
A 89482-E7	4.0	
oneer 2691	eig a v	医二硫甲酚酚 医环形二氏管 电电阻电阻
92944RCX	2.7	Regional Control of the Control of t
931839A5	1.3	0.04
\ 81404-E56	1.3	0.07
X 88130-LE5	0	0.01
Stuckey	0	o
D 0.1	19.9	0.57

^{*} indicates mean is significantly greater than zero (P < 0.1; LSD test).

Susceptible in previous tests.

Resistant in previous tests.

Hessian infestations in entries of the State Wheat Variety Trial at Plains, GA in spring of 1997.

P-4	% Infested	Immatures
Entry CA 97467 F24	stems	per stem
• GA 87467-E24	16.0 *	0.28 *
Kucharek A28	16.0 *	0.21 *
• Featherstone 520	16.0 *	0.21 *
• Clemens	13.3 *	0.29 *
Agratech FFR 522W	12.0 *	0.32 *
• Pioneer 2643	12.0 *	0.16 *
• Coker 9803	12.0 *	0.13 *
• Mason	6.7	0.07
■ Madison	6.7	0.09
Pioneer 2580	6.7	0.08
Coker L910097	6.7	0.09
Morey	5.3	0.09
Clemson 201	4.0	0.04
Jackson	4.0	0.04
Jaypee	4.0	0.04
Agripro 92D+4413	4.0	0.13
Hickory	2.7	0.03
Florida 304	2.7	0.03
Pioneer 2691	2.7	0.04
GA 87105-E43	2.7	0.03
Hunter	2.7	0.08
Coker 9663	2.7	0.04
GA Gore	1.3	0.01
GA Dozier	1.3	0.01
Agratech FFR 502W	1.3	0.03
GA 871339-E18	1.3	0.01
Pioneer 2684	1.3	0.01
FL 92944RCX	1.3	0.01
Clark	0	0
Coker 9134	0	0
Coker 9835	0 🚕	Salvania XO
Agratech EX096W	0	
Pioneer 2628	. 0	Ö
GA Stuckey		
GA Fleming (90078-I)	0	0.000
FL 931839A5	Ö	0
GA 89482-E7	Ö	0
GA 881186-E48	Ö	0
GA 881404-E56	. 0	0
GA 881130-LE5	. 0	. 0
GA 891283-LE18	0	
VA 93-52-60	0	0
LSD 0.1	7.9	0
LSD 0.05	7. 9 9.4	0.13

<sup>indicates mean is significantly greater than zero (P < 0.1; LSD test).
Susceptible in previous trials.</sup>

[■] Resistant in previous trials.

Table 13
Hessian fly infestations in entries of the State Wheat Variety
Test at Plains, GA in 1995/1996.

Debase	\$ Infested	Immatures				
Entry	Plants	per plant				
●Jackson						
Featherstone 520	34.5*	0.65*				
ABI 9ZD-4413	32.8*	0.83*				
•Pioneer 2643	32.6*	0.62*				
Clemens	28.3*	0.37*				
•NK Coker 9803	25.4*	0.44*				
Mason	25.1*	0.36*				
•GA 87467-E24	23.7*	0.55*				
•Savannah	23.6*	0.37*				
ARK 26158-4	22.9*	0.37*				
GA 88509-E31	19.5*	0.30*				
Pioneer 2691	19.2*	0.25*				
•Clemson 201	17.4*	0.22*				
●FL 85238-G585	17.2*	0.22*				
NK Coker X 900819	16.4*	0.24*				
•Clark	14.7*	0.26*				
	13.6*	0.21*				
T 8725-E11	11.7*	0.16				
◆GA 90078-I	10.8*	0.17				
♦Hickory	9.4	0.12				
♦Pioneer 2580	8.9	0.16				
GA 88151-L19	6.4	0.10				
♦Madison	4.0	0.06				
Pioneer 2684	3.7	0.04				
GA 87105-E43	3.2	0.05				
GA 88355-L16	2.9	0.06				
♥GA-Gore	1.3	0.01				
Florida 304	1.1	0.01				
VA 93-52-60	0.9	0.01				
Pioneer 2628	0.8	0.01				
GA Andy	0.6	0.01				
GA 89253-E39	0.6	0.01				
NK Coker 9134	AD ON THE ST	. ()				
Stacy	O The second of	0				
GA 871339-E18	0	0				
GA 87436-E14	0	₂₁ 0				
GA Dozier	120 00.39					
NK Coker 9835		0				
GA Stuckey	o	0				
GA 88129-E11	o	0				
Morey	· 0	0				
SD (0.1)	10.3	0.18				
SD (0.05)	12.3	0.21				

Trial consists of 3 ft rows arranged in a randomized complete block design with 3 replicates; counts are from 1 ft of row per plot, collected on Feb. 14, 1996 when plants were full tillering.

^{*} significantly greater than zero (P<0.1).

These entries have been susceptible in previous trials.

[◆] These entries have been resistant in previous trials.

1996-97 Uniform Southern Soft Red Wheat Performance Nursery

\$				
Contents	<u>Index</u>	Rust News	Mail Lists	<u>Home</u>

Seedling reaction of entries of the 1996-97 Uniform Southern Soft Red Wheat Performance Nursery to selected isolates of *Puccinia recondita* f. sp. *tritici* (comments or questions? Send a message to <u>David L.Long</u>)

Reactions produced by Prt race

No. Cultivar or Line	PLMQ	МСЛ	TCDL	LBBQ	SBJB	TCBG	TLGG	СВТВ	MBRL	PNM	Postulated LLr genes*:
1 FL 302	3	3	3	,	•				3	3	3,10
2 Coker 9835			;	;1c	•	,	3	•	;		2a,9,11
3 Pioneer 2643	;-3		;	,			:	3;	;		+
4 SC 900237	;	• ;	,	•	,	,		;		;	++
5 L910097		;	3	3;	,	•	;3	;	;2	:	2a,10,+
6 MO 94-317	,		;1 c	į	,	,1c	;	;2	3-;	:	. +
7 SC 910031	-	3-;	;1c	;1c	•	;1c	;	,1c	;lc		+
8 GA 87467	esta anticoloria. Regionale										+ 8
9 GA 871339	-	;-3	,	;1c		,	,1c	•		3-,	+
10 GA 90078		;1c	;-3	3	3	;-3	,	3		;1c	+
11 LA 85422C13142	;	;	;1c	;1c	;1c	,	;	;1c3	3;	;1c1	+
12 TX 91D6999	•	;-3	;-3		•	;	;1c	;-3			+
13 VA 94-52-68		;	,1c	್ರ,2c ျ	,lc	,	,	;1 c	;1c	,	+
14 VA 94-54-479	•	;	;-3	esserie	;-3			;1c-3	;1c	;1c	+
15 VA 94-54-549	-	;		;1c	3	•		,	;1c2		2a,11,17,+
16 AR 494B-2-2	3	3	33	(1ેટ3) ∀	•	;-3	;3	•	3	3	10
17 AR 584A-3-2	•	;	. ;	11. , 150	(5,4)		;1 c	,		3	+
18 FL 92944RCX		;	;	;1c					;lc		+
19 FL 931339AS	•	;	;	,		•	,	,			+
20 FL 92944BX	;	,	,	;	· •	;	;	;	;1c	,	+
21 A93-6061	-	;	;	. ;	3	;1c	3;	;	,	,	2a,11,17,+
22 A93-6227	;	,	;		3;	3-;	,	,	•	;1c	+
23 A93-7162	;	•	;	;1c	•	;		;	•	;1c	+
24 L920738	;1c	,	;	3-;	,	;	,	,	,	,	+
25 L920024	;	•	3	3;	,	,	,	,	;1c	,	2a,10,+
26 LA8889B213112	•	•	,	;	;	,	;			;1c	+

27 LA					0.1						
8529B35112B1	,	,	,	,	;21c	;1c	,	;	;	,	+
28 LA87167D8102	;			;lc	•		;	:	· ;	;	+
29 TX 92D7702	3	3	3	3	3;	3;	3	3	3	3	0
30 TX92D8102	;1c	,	;	,	3	<i>,</i>		,	,		2a,11,17,+
31 NCV93-1007	•	;	,	,	,	,	•	,			+
32 NCV93-612	;	;	3	;lc	,	,lc			· ,		2a,10,+
33 TX18NT	;1c	;	,	;1 c	;2c	;-3	;3	;1 c 3	;3	;-3	+

^{*}Single genes tested = 1, 2a, 2c, 3, 3ka, 9, 10, 11, 16, 17, 18, 24, 26, 30

Virulence Formula:

PLMQ = 1, 2c, 3, 3ka, 9, 10, 18, 30

MCJL = 1, 3, 10, 11, 17, 26

TCDL = 1, 2a, 2c, 3, 10, 17, 26

TBBQ = 1, 2a, 2c, 3, 9, 11, 18

CBTB = 3, 3ka, 11, 17, 30

MBRL = 1, 3, 3ka, 10, 11, 30

SBJB = 1, 2a, 2c, 11, 17

PNML = 1, 2c, 3, 3ka, 9, 10, 24, 30

<u>Introduction - What's New - Research - Publications - Databases - Barberry Testing</u>
<u>Rust Bulletins - Rust Losses - Image Gallery - Other Resources - Home</u>

CO (MW SE

NEDVATOR HARD VELEMBERH.

^{**0 =} no gene(s) detected with these Lr combinations; + = Lr gene(s) present but unable to identify with these Lr virulence combinations

							Τ	Ť	1		· -		T		
LAB		ENTRY	MILLIN G			BAKIN G			COMBIN ED		MICR O	SOFT		FLOU R	FLOU
NO.			QUALI TY			QUALI TY			QUALIT Y		T.W.	EQUI V		YIEL D	PROT
			SCOR E			SCOR E			SCORE		LB/B U				
****		CTANGAGO		-	╀				ļ	_			-	<u> </u>	
	4	STANDARD	100.0	+	\perp	100.0		_	100.0		58.4	62.2	+	68.6	7.5
1801		FL 302	101.8		\vdash	95.4			95.4		56.4 *	59.6	\perp	69.6	8.4
1802		Coker 9835	100.0	1-	-	100.0	1	\perp	100.0		58.4	62.2	╀-	68.6	7.5
1803		Pioneer 2643	94.9	+	\vdash	100.7	 	-	94.9	\vdash	57.5	57.8 *	\vdash	68.1	8.5
1804 1805		SC 900237	79.9	_	\vdash	84.7		\perp	79.9		59.1	57.0 *	-	64.3 Q	8.7
1806		L910097	96.4	+	-	94.3		╀	94.3		63.8	57.2 *	-	68.1	9.2
1807		MO 94-317	104.7	+		107.5		╀	104.7		57.8	62.3	-	70.3	8.2
1808		SC 910031 GA 87467	87.7			94.4			87.7		57.1 *	51.1 Q	-	67.4 *	9.1
1809			91.4			102.4		100	91.4		58.4	56.6 *	100	67.3	8.2
1810		GA 871339 GA 90078	90.4		\vdash	83.3		╁	83.3		57.0 *	56.5 *	╀	67.2 *	8.4
1811			91.0		\vdash	74.8		+	74.8		58.9	53.9 Q	+	67.5 *	8.2
1812		LA 85422-C13-1-4-2 TX 91D6999	100.9		Н	97.7		╀	97.7	\rightarrow	59.0	55.0 C	+	70.2	7.9
1813		VA 94-52-68	77.9	 	Н	61.2		+-	61.2		57.9	50.1 Q	-	65.0 Q	9.3
1814		VA 94-52-66 VA 94-54-479	95.0		Н	85.5		+	85.5		59.3	56.4 *	H	68.1	9.1
1815		VA 94-54-549	88.9		Н	72.1		+	72.1		58.9	53.2 Q	-	67.2 *	8.0
1816		AR 494B-2-2	92.0	_		87.1		┼┈	87.1		59.3	56.4 *	+-	67.4 *	9.0
1817		AR 584A-3-2	95.8			89.9		+	89.9		58.0	56.1 *	\vdash	68.5	8.5
1818		FL92944RCX	100.3		\vdash	100.1		-	100.1		59.1	54.0 Q	-	69.9	8.9
1819		FL931339AS	99.2		Н	105.3		╄	99.2		56.7 *	61.7	├	68.7	8.3
1820		FL92944BX	76.9 100.4		\vdash	62.1		╁	62.1	-	58.5	47.0 Q		65.1 Q	9.8
1821		A93-6061	98.0			97.6 100.1		-	97.6 98.0		55.7 Q	1	\vdash	69.3 68.7	8.7
1822		A93-6227	101.7	_		94.6		-			55.4 Q	60.3 55.0 Q	├		8.6
1823		A93*7162	89.9			70.2		╁	94.6 70.2		60.6	52.5 Q	+-	70.4 67.3 *	9.4
1824		L920738	98.1		\vdash	103.8		-	98.1		58.0	59.8	╁	68.5	9.3
1825		L920024	92.0			87.9		-	87.9		59.9	55.0 Q		67.5 *	9.2
1826		LA8889-B-2-1-3-1-1-2	99.7		-	78.3			78.3		60.6	52.1 Q		69.8	9.2
1827		LA8529-B-3-5-11-2-B-1	93.3	-	\vdash	86.8		-	86.8		58.8	54.7 Q	_	68.0	9.2
1828		LA87167-D8-10-2	97.9	$\overline{}$	H	93.5		+	93.5		59.7	51.4 Q	+	69.5	8.7
1829		TX92D7702	95.9			108.0	• • • • •	+	95.9	$\overline{}$	54.0 Q		 	68.0	8.2
1830		TX92D8102	87.9		H	38.4			38.4		57.5	44.3 Q		68.3	8.9
1831		NCV93-1007	102.8		\forall	105.2		H	102.8		56.8 *	59.7		70.5	
1832		NCV93-612				103.2		Н	102.8		60.0	59.7		70.5	8.6
1833		TX18NT	96.1		البياة	90.9		Н	90.9	_	58.7	59.0		68.0	8.7

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in 1974 (5 U.S.C. 552a) and the Paperwor	
EXHIBIT E	Application is required in order to de	
STATEMENT OF THE BASIS OF OWNERSHIP	certificate is to be issued (7 U.S.C. 24 until certificate is issued (7 U.S.C. 2426)	
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
University of Georgia Research Foundation	GA87467E24	518W
Florida Agricultural Experiment Station		
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
Boyd Graduate Studies Research Center Room 630	706-542-6512	706-542-3837
Athens, GA 30602-7411	7. PVPO NUMBER	
	98	300327
8. Does the applicant own all rights to the variety? Mark an "X" in appropri	iale block. If no, please explain.	YES NO
 Is the applicant (individual or company) a U.S. national or U.S. based of If no, give name of country 	ompany?	YES NO
10. Is the applicant the original owner?	O If no, please answer one of the fo	ollowing:
a. If original rights to variety were owned by individual(s), is (are) the or	Callenoiten 2 I s (abanua Iraini	
	· _	0
b. If original rights to variety were owned by a company(ies), is(are) the		/ • ·
∑ YES □ N	If no, give name of country	
11. Additional explanation on ownership (if needed, use reverse for extra s	pace):	•
SEE ATTACHMENT		· · · · · · · · · · · · · · · · · · ·
PLEASE NOTE:		
Plant variety protection can be afforded only to owners (not licensees) who meet o	ne of the following criteria:	
1. If the rights to the variety are owned by the original breeder, that person must be which affords similar protection to nationals of the U.S. for the same genus and	e a U.S. national, national of a UPOV members.	per country, or national of a country
If the rights to the variety are owned by the company which employed the originember country, or owned by nationals of a country which affords similar prot	inal breeder(s), the company must be U.S. be ection to nationals of the U.S. for the same	used, owned by nationals of a UPOV genus and species.
3. If the applicant is an owner who is not the original owner, both the original own		
The original breeder/owner may be the individual or company who directed final l		
According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collisis information collection is 0581-0055. The time required to compete this information collection is 0581-0055. The time required to compete this information collecting exciting existing data sources, gathering and maintaining the data needed, and completing and	JOU & SAMPAGE IN SAME IN UNIVER HE LESIVE.	critrol number. The valid OMB control number for se, including the time for reviewing instructions,
The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative USDA's TARGET Center at 202-720-2600 (voice and TDD).	of more color national prioring sext religions age, distab	ility, political beliefs, and marital or familial status. braille, targe print, audiotape, etc.) should contact
To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Was	hington, D.C. 20250, or call 1-800-245-6340 (voic	a) or (202) 720-1127 (TOD). USDA is an equal

employment opportunity employer.

STD-470-E (07-97) (Destroy previous editions).
Electronic version designed using WordPerfect InForms by USDA-AMS-IMB.

Exhibit E Statement of Applicant's Ownership 518W

The variety for which plant variety protection is hereby sought is owned jointly by the University of Georgia Research Foundation, Inc. (UGARF) and the Florida Agricultural Experiment Station, University of Florida (FAES).

Ownership by UGARF in the variety for which plant variety protection is hereby sought is based on the Patent Policy approved by the Board of Regents of the University System of Georgia on June 9, 1982, in which the Board of Regents assigned to the University of Georgia Research Foundation, Inc. all rights in intellectual property developed or created by employees at The University of Georgia, one of the universities of the University System of Georgia. Rights of novel plant varieties developed at The University of Georgia, including '518W', are covered by said Patent Policy. As employees of The University of Georgia, Jerry W. Johnson, Barry Cunfer, and G. David Buntin, pursuant to said Patent Policy, have assigned their right sin '518W' to the University of Georgia Research Foundation, Inc.

Ron Barnett and Paul Pfahler are employees of the Florida Agricultural Experiment Station, the University of Florida.